

## EXPRESS MAIL CERTIFICATE

I hereby certify that this paper or fee is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 C.F.R. § 1.10 on the date indicated above and is addressed to the Assistant Commissioner for Patents, Washington, D.C. 20231

Typed or Printed Name	Margaret Pierce	Express Mail No.	EL 563 388 109 US
Signature	<i>M. Pierce</i>	Date	2/20/01

Form PTO-1390  
(REV 10-94)

U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE

TRANSMITTAL LETTER TO THE UNITED STATES  
DESIGNATED/ELECTED OFFICE (DO/EO/US)  
CONCERNING A FILING UNDER 35 U.S.C. 371

ATTORNEY'S DOCKET NUMBER  
SHIM-008U.S. APPLICATION NO.  
To Be Assigned

09/763286

INTERNATIONAL APPLICATION NO. PCT/JP99/04251	INTERNATIONAL FILING DATE 23 August 1999	PRIORITY DATE CLAIMED 24 August 1998
---	---	---

TITLE OF INVENTION  
CARBONYL-STRESS IMPROVING AGENT AND PERITONEAL DIALYSATE

APPLICANT(S) FOR DO/EO/US  
Toshio Miyata

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. ☒ This express request to begin examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and the PCT Articles 22 and 39(1).
4. ☒ A proper Demand for International Preliminary Examination was made by the 19<sup>th</sup> month from the earliest claimed priority date.
5. ☒ A copy of the International Application as filed (35 U.S.C. 371(c)(2)):
  - a. ☐ is transmitted herewith (required only if not transmitted by the International Bureau).
  - b. ☒ has been transmitted by the International Bureau. (**However, a courtesy copy is attached hereto.**)
  - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US).
6. ☐ A translation of the International Application into English (35 U.S.C. 371(c)(2)).
7. ☒ Amendments to the claims of the International Application under PCT Article 19 (37 U.S.C. 371(c)(3)):
  - a. ☐ are transmitted herewith (required only if not transmitted by the International Bureau).
  - b. ☐ have been transmitted by the International Bureau.
  - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
  - d. ☒ have not been made and will not be made.
8. ☐ A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
9. ☒ An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)). (**UNSIGNED**)
10. ☐ A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).

**Items 11 to 16 below concern other document(s) or information included:**

11. ☐ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
12. ☐ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
13. ☐ A **FIRST** preliminary amendment.  
☐ A **SECOND** or **SUBSEQUENT** preliminary amendment.
14. ☐ A substitute specification.
15. ☐ A change of power of attorney and/or address letter.
16. ☒ Other items or information: Certification Regarding Sequence Listing (1 page); Paper Copy of Sequence Listing (2 pages); Diskette containing sequence listing

U.S. APPLICATION NO. 11 (known, see 37 CFR 1.20) To Be Assigned: <b>09/763286</b>		INTERNATIONAL APPLICATION NO. PCT/JP99/04521		ATTORNEY'S DOCKET NUMBER SHIM-008	
17. <input checked="" type="checkbox"/> The following fees are submitted: <b>Basic National Fee (37 CFR 1.492(a)(1)-(5)):</b>  Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO and international search report not prepared by the EPO or JPO \$970.00  International preliminary examination fee (37 CFR 1.482) not paid to USPTO but international search report prepared by the EPO or JPO \$840.00  International preliminary examination fee (37 CFR 1.482) not paid to USPTO but international search fee (37 CFR 1.445(a)(2)) paid to USPTO \$690.00  International preliminary examination fee (37 CFR 1.482) paid to USPTO but all claims did not satisfy provisions of PCT Article 33(1)-(4) \$670.00  International preliminary examination fee (37 CFR 1.482) paid to USPTO and all claims satisfied provisions of PCT Article 33(1)-(4) \$ 96.00					
<b>ENTER APPROPRIATE BASIC FEE AMOUNT =</b>				\$ 840.00	
Surcharge of <b>\$130.00</b> for furnishing the oath or declaration later than _____ 20 _____ 30 months from the earliest claimed priority date (37 CFR 1.492(e)).				\$ 0.00	
CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE		
Total Claims	16 - 20 =	0	X \$18.00	\$ 0.00	
Independent Claims	2 - 3 =	0	X \$78.00	\$ 0.00	
MULTIPLE DEPENDENT CLAIM(S) (if applicable)			+ \$260.00	\$ 260.00	
<b>TOTAL OF ABOVE CALCULATIONS =</b>				\$ 1100.00	
Applicant claims small entity status. See 37 CFR 1.27. Reduction of 1/2 for filing by small entity.				\$ 550.00	
<b>SUBTOTAL =</b>				\$ 550.00	
Processing Fee of <b>\$130.00</b> for furnishing the English translation later than _____ 20 _____ 30 months from the earliest claimed priority date (37 CFR 1.492(f)).				\$ 0.00	
<b>TOTAL NATIONAL FEE =</b>				\$ 550.00	
Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 per property.				\$ 0.00	
<b>TOTAL FEES ENCLOSED =</b>				\$ 550.00	
				<b>Amount to be: refunded</b>	\$
				<b>charged</b>	\$

a. .... A check in the amount of \$ \* to cover the above fees is included.

b. ... ☒ Please charge my Deposit Account No. 50-0815 in the amount of \$ 550.00 to cover the above fees. A duplicate copy of this sheet is enclosed.

c. ... ☒ The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. 50-0815. A duplicate copy of this sheet is enclosed.

**NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.**

SEND ALL CORRESPONDENCE TO

BOZICEVIC, FIELD & FRANCIS LLP  
200 Middlefield Road, Suite 200  
Menlo Park, CA 94025  
(650) 327-3400 Telephone  
(650) 327-3231 Facsimile  
.....

SIGNATURE

Karl Bozicevic

NAME

28,807

REGISTRATION NO.

## DESCRIPTION

## CARBONYL-STRESS IMPROVING AGENT AND PERITONEAL DIALYSATE

5 Technical Field

The present invention relates to a peritoneal dialysate used for treating patients with renal failure.

Background Art

10 Two types of dialysis, hemodialysis and peritoneal dialysis, are used to treat patients with chronic renal failure. Peritoneal dialysis is a method in which the dialysate is allowed to dwell in the peritoneal cavity for a certain period of time, thereby facilitating the excretion of waste products out of the body into the dialysate through the peritoneum. The dialysate is then recovered. Peritoneal dialysis is subdivided into intermittent peritoneal dialysis (IPD) and continuous ambulatory peritoneal dialysis (CAPD). CAPD is a method that incorporates the merits of the IPD method in which the fluid exchange is carried out about four times a day by lengthening the dwelling time of the perfusate in the peritoneal cavity.

20 Peritoneal dialysis has advantages such as being convenient and less time-consuming. However, it is known that long-term treatment with peritoneal dialysis can progressively lower the ability of water removal, and can result in abdominal protein denaturation and hardening, peritoneal fusion, and such abnormalities.

A part of the cause is thought to be glucose present in the peritoneal dialysate. Many types of peritoneal dialysates used today contain glucose as an osmoregulatory agent. Glucose is unstable to heat, and a part thereof is degraded during heat sterilization. As a result, highly reactive carbonyl compounds capable of modifying proteins may generate as degradation products. Such degradation products may also generate and accumulate in a glucose-containing peritoneal dialysate even during storage that follows sterilization.

35 Generally, glucose is apt to degrade at a nearly neutral or alkaline pH, and therefore, acidic buffers (pH 5.0-5.4) are selected to maintain the stability of glucose in ordinary peritoneal dialysates. However,

such acidic buffers carry risks such as suppressing immunological defense mechanisms of peritoneal macrophages, causing the onset of peritonitis due to bacterial infection, and being cytotoxic to peritoneal mesothelial cells. To overcome such mutually  
5 contradictory problems, there was a desperate need to prevent generation of carbonyl compounds resulting from the degradation of glucose within peritoneal dialysates around a neutral pH, or eliminate such compounds.

On the other hand, a peritoneal dialysate formulated with a high  
10 concentration of glucose can modify proteins, and therefore, such dialysates are unfavorable for the peritoneum. From such a viewpoint, some peritoneal dialysates have been developed by utilizing glucose polymers that generate few degradation products (Unexamined Published Japanese Patent Application (JP-A) NO. Hei 10-94598; Wilkie, M.E.  
15 et al., Perit. Dial. Int., 17: S47-50 (1997)).

From the same viewpoint, other compounds have been proposed in place of glucose as osmoregulatory agents used in peritoneal dialysates. These include, cyclodextrin (JP-A Hei 8-71146), disaccharide (JP-A Hei 8-131541), and amino acids (Faller, B. et al.,  
20 Kidney Int., 50 (suppl.56), S81-85 (1996)). A peritoneal dialysate having cysteine as an additive to prevent the degradation of glucose has also been disclosed (JP-A Hei 5-105633).

These methods aim to improve inconveniences caused by the high concentration of glucose within the peritoneal dialysate.

It has been reported that, irrespective of the presence or absence  
25 of hyperglycemia, large amounts of highly reactive carbonyl compounds and AGE (advanced glycation end products) are accumulated in the blood and tissues of patients with chronic renal failure (Miyata, T. et al., Kidney Int., 51:1170-1181 (1997); Miyata, T. et al., J. Am. Soc.  
30 Nephrol., 7: 1198-1206 (1996); Miyata, T. et al., Kidney Int. 54:1290-1295 (1998); Miyata, T. et al., J. Am. Soc. Nephrol. 9:2349-2356 (1998)). Renal failure often accompanies conditions having an overload of carbonyl compounds (carbonyl stress). This carbonyl stress results from non-enzymatic biochemical reactions to generate  
35 carbonyl compounds from sugars and lipids, which is thought to lead to enhanced protein modifications (Miyata, T. et al., Kidney Int.

55:389-399 (1999)). Carbonyl stress not only alters the architecture of matrix proteins such as collagen and fibronectin, but also participates in the enhancement of peritoneal permeability and the onset of inflammation due to the physiological activities of carbonyl compounds towards a variety of cells.

In peritoneal dialysis, waste products in the blood are excreted into the peritoneal dialysate through the peritoneum. A hyperosmotic peritoneal dialysate dwelling within the peritoneal cavity collects highly reactive carbonyl compounds accumulated in the blood of renal failure patients through the peritoneum into itself. Thus, the carbonyl-compound concentration in the peritoneal dialysate elevates, resulting in a carbonyl-stress state. This is thought to cause carbonyl modification of proteins in the peritoneal cavity and as a consequence, the peritoneal functions are suppressed to advance peritoneal sclerosis.

Immunohistochemical examination of the endothelium and mesothelium, has demonstrated that the carbonyl-stress state is caused by glucose present in the peritoneal cavity in peritoneal-dialysis patients (Yamada, K. et al., Clin. Nephrol., 42: 354-361 (1994); Nakayama, M. et al., Kidney Int., 51: 182-186 (1997); Miyata, T. et al., J. Am. Soc. Nephrol. in press; Combet, S. et al., J. Am. Soc. Nephrol. in press; Inagi, R. et al., J. Am. Soc. Nephrol. in press).

#### Disclosure of the invention

An objective of the present invention is to provide a method for reducing damage from carbonyl compounds during peritoneal dialysis, specifically for improving the carbonyl-stress state, as well as to provide a dialysate and pharmaceutical agent required for this method. The "carbonyl compound" in the present invention means a carbonyl compound derived from patients undergoing peritoneal-dialysis, a carbonyl compound generated in the peritoneal dialysate during manufacturing or storing, and a carbonyl compound generated in the peritoneal cavity during peritoneal dialysis. The present invention aims to minimize as much as possible the damages from these carbonyl compounds on the patients undergoing dialysis.

The present inventors have discovered that the highly reactive

carbonyl compounds present in the peritoneal dialysate in peritoneal dialysis patients are not merely those present in the original peritoneal dialysate that have been infused into the peritoneal cavity. Specifically, the amount of glucose-excluded carbonyl compounds in the peritoneal dialysate recovered from peritoneal-dialysis patients is five times more than the amount before dialysis, and thus, the majority of the carbonyl compounds is assumed to derive from blood (Fig. 1). Therefore, it has been revealed that the carbonyl compounds present in the peritoneal dialysate in the peritoneal cavity, include not only carbonyl compounds generated in the process of heat sterilization and those that accumulate during storage, but also those that are derived from blood and those that generate and accumulate within the peritoneal cavity during dialysis, the latter two being not negligible. Immunohistochemical examinations of peritoneal tissue from peritoneal-dialysis patients have revealed the presence of carbonyl-modified proteins (Fig. 2). Accordingly, if it is possible to remove carbonyl compounds transferred from the blood into the peritoneal cavity during peritoneal dialysis, the carbonyl-stress state would also be effectively improved.

The present inventors thought that renal failure should often accompany conditions in which *in vivo* protein modification was enhanced and that, when glucose was continuously infused at a high concentration into the peritoneal cavity by a method such as peritoneal dialysis, intraperitoneal proteins would be increasingly susceptible to non-enzymatic modifications due to the peritoneal dialysate in which peritoneal cavity-derived carbonyl compounds were accumulated (Fig. 7).

Based on such a background, the present inventors have discovered that a carbonyl compound-trapping agent is effective in preparing a dialysate capable of reducing carbonyl compounds of peritoneal dialysate origin, and thus, completed the present invention. Focusing mainly on carbonyl compounds that accumulate within blood, the present inventors have further found that pharmaceutical agents that prevent protein modifications caused by carbonyl stress and such peritoneal dialysis-associated complications, are useful, thereby further completing the present invention.

Specifically, the present invention relates to a carbonyl-stress improving agent, and a peritoneal dialysate based thereon, as well as a pharmaceutical agent. The present invention relates more specifically to:

5 (1) an agent for improving intraperitoneal carbonyl-stress state during peritoneal dialysis, comprising a carbonyl compound-trapping agent as an active ingredient;

(2) the agent of (1), wherein the carbonyl compound-trapping agent is immobilized on an insoluble carrier;

10 (3) the agent of (1), wherein the carbonyl compound-trapping agent is to be mixed with a peritoneal dialysate;

(4) the agent of any one of (1) to (3), wherein the carbonyl compound-trapping agent is selected from the group consisting of aminoguanidine, pyridoxamine, hydrazine, SH group containing compound, and derivatives of these;

15 (5) the agent of any one of (1) to (3), wherein the carbonyl compound-trapping agent is an agent inhibiting Maillard reaction;

(6) the agent of (1), wherein the carbonyl compound-trapping agent is a compound insoluble in peritoneal dialysates and capable of adsorbing carbonyl compounds;

20 (7) a cartridge used for trapping carbonyl compounds within peritoneal dialysates, wherein the cartridge is filled with the carbonyl compound-trapping agent(s) of (2) and/or (6);

25 (8) a method for preparing a peritoneal dialysate having a reduced carbonyl compound content, the method comprising passing the peritoneal dialysate through the cartridge of (7);

(9) a method for preparing a peritoneal dialysate having a reduced carbonyl compound content, the method comprising:

30 (a) contacting the peritoneal dialysate with the carbonyl compound-trapping agent(s) of (2) and/or (6) and

(b) separating the peritoneal dialysate from the carbonyl compound-trapping agent;

(10) a peritoneal dialysate comprising a carbonyl compound-trapping agent;

35 (11) the peritoneal dialysate of (10), wherein the peritoneal dialysate further comprises a reducing sugar, and is placed in a

09763286, 082001

container comprising a first compartment and a second compartment so that the first compartment contains the reducing sugar and the second compartment contains the carbonyl compound-trapping agent; and

- 5 (12) the peritoneal dialysate of (10), wherein the carbonyl compound-trapping agent is to be administered into the intraperitoneal cavity.

The present invention further relates to the use of carbonyl compound-trapping agent for improving intraperitoneal carbonyl-stress state. The present invention also relates to the use of carbonyl compound-trapping agent for the peritoneal dialysis treatment. Furthermore, the present invention relates to the use of carbonyl compound-trapping agent for producing an agent improving carbonyl stress.

15 In the present invention, the carbonyl compounds to be trapped include, for example, carbonyl compounds generated in the processes of production and storage of a peritoneal dialysate. As described above, carbonyl compounds can always be generated in a peritoneal dialysate containing a high concentration of glucose as an osmoregulatory agent. Such carbonyl compounds include, for example, the following compounds (Richard, J. U. et al., Fund. Appl. Toxic., 4: 843-853 (1984)):

- 3-deoxyglucosone
- 5-hydroxymethylfurfural (abbreviated hereafter "5-HMF")
- 25 • formaldehyde
- acetaldehyde
- glyoxal
- methylglyoxal
- levulinic acid
- 30 • furfural
- arabinose

In the present invention, a carbonyl compound-trapping agent is used throughout the dialysis, thereby achieving the removal of carbonyl compounds, as listed below, which are accumulated in the blood of a patient with renal failure and are transferred into the peritoneal cavity following peritoneal dialysis, as well as the removal of carbonyl



compounds generated in a peritoneal dialysate during its production process and storage process.

Carbonyl compounds derived from ascorbic acid:

- dehydroascorbic acid

Carbonyl compounds derived from carbohydrate, lipid, or amino acid:

- glyoxal
- methylglyoxal
- 3-deoxyglucosone
- hydroxynonenal
- malondialdehyde
- acrolein

A preferred carbonyl compound-trapping agent in the present invention is one capable of completely inhibiting or reducing the protein-modification activity of all these carbonyl compounds through a chemical reaction or adsorption; however the carbonyl compound-trapping agent also includes an agent effective merely for major carbonyl compounds among these. For example, methylglyoxal is believed to have a relatively high reactivity among carbonyl compounds (see Thornalley, R.J., Endocrinol. Metab. 3 (1996) 149-166; Inagi, R. et al., J. Am. Soc. Nephrol. in press; and Example 3), and therefore it is of great pathophysiological significance to inhibit the activity of methylglyoxal. Consequently, a compound effective for methylglyoxal can be a preferred carbonyl compound-trapping agent of the present invention. Specifically, as indicated in Examples, compounds such as activated charcoal, guanidine, aminoguanidine, biguanide, cysteine, and albumin are particularly effective carbonyl compound-trapping agents for methylglyoxal.

Although some carbohydrates used as osmoregulatory agents have been reported to be more stable than glucose, it is difficult to completely inhibit the generation of carbonyl compounds from these carbohydrates during heat sterilization and storage processes. Thus, the use of the carbonyl compound-trapping agent is meaningful in situations where carbohydrates other than glucose are used as osmoregulatory agents. Carbohydrates other than glucose usable as osmoregulatory agents in peritoneal dialysis, include, trehalose (JP-A

Hei 7-323084), hydrolyzed starch (JP-A Hei 8-85701), maltitol and lactitol (JP-A Hei 8-131541), as well as non-reducing oligosaccharides and non-reducing polysaccharides (JP-A Hei 10-94598).

Carbonyl compound-trapping agents of the present invention include compounds capable of inhibiting or reducing the damage due to carbonyl compounds towards dialysis patients by a chemical reaction or adsorption, and which per se is safe for the patients. Such compounds include those listed below. The inventive carbonyl compound-trapping agent can be used alone or in combination of two or more types of compounds.

• aminoguanidine (Foote, E. F. et al., Am. J. Kidney Dis., 25: 420-425 (1995))

•  $\pm$  2-isopropylidenehydrazono-4-oxo-thiazolidin-5-yl-acetanilide (OPB-9195; S. Nakamura, 1997, Diabetes 46:895-899)

Further, the carbonyl compound-trapping agent includes, for example, the following compounds or derivatives thereof that are capable of functioning as a carbonyl compound-trapping agent. The "derivatives" indicate compounds having an atomic or molecular substitution(s) at any position as compared with the parental compound.

(1) guanidine derivatives such as methylguanidine (JP-A Sho 62-142114; JP-A Sho 62-249908; JP-A Hei 1-56614; JP-A Hei 1-83059; JP-A Hei 2-156; JP-A Hei 2-765; JP-A Hei 2-42053; JP-A Hei 6-9380; Published Japanese Translation of International Publication 5-505189), etc.

(2) hydrazine derivatives such as sulfonylhydrazine, etc.

(3) 5-membered heterocyclic compounds having 2 nitrogen atoms, such as pyrazolone (JP-A Hei 6-287179), pyrazoline (JP-A Hei 10-167965), pyrazole (JP-A Hei 6-192089; JP-A Hei 6-298737; JP-A Hei 6-298738), imidazolysine (JP-A Hei 5-201993; JP-A Hei 6-135968; JP-A Hei 7-133264; JP-A Hei 10-182460), hydantoin (JP-A Hei 6-135968), etc.

(4) 5-membered heterocyclic compounds having 3 nitrogen atoms, such as triazole (JP-A Hei 6-192089), etc.

(5) 5-membered heterocyclic compounds having a nitrogen atom and a sulfur atom, such as thiazoline (JP-A Hei 10-167965), thiazole (JP-A Hei 4-9375; JP-A Hei 9-59258), thiazoline (JP-A Hei 5-201993; JP-A

Hei 3-261772; JP-A Hei 7-133264; JP-A Hei 8-157473), etc.

(6) 5-membered heterocyclic compounds having a nitrogen atom and an oxygen atom, such as oxazole (JP-A Hei 9-59258), etc.

5 (7) nitrogen-containing 6-membered heterocyclic compounds such as pyridine (JP-A Hei 10-158244; JP-A Hei 10-175954), pyrimidine (Published Japanese Translation of International Publication 7-500811), etc.

10 (8) nitrogen-containing condensed heterocyclic compounds such as indazole (JP-A Hei 6-287180), benzimidazole (JP-A Hei 6-305964), quinoline (JP-A Hei 3-161441), etc.

(9) sulfur- and nitrogen-containing condensed heterocyclic compounds such as benzothiazole (JP-A Hei 6-305964), etc.

(10) sulfur-containing condensed heterocyclic compound such as benzothiophene (JP-A Hei 7-196498), etc.

15 (11) oxygen-containing condensed heterocyclic compounds such as benzopyran (JP-A Hei 3-204874; JP-A Hei 4-308586), etc.

(12) nitrogenous compounds such as carbazoyl (JP-A Hei 2-156; JP-A Hei 2-753), carbazic acid (JP-A Hei 2-167264), hydrazine (JP-A Hei 3-148220), etc.

20 (13) quinones such as benzoquinone (JP-A Hei 9-315960), hydroquinone (JP-A Hei 5-9114), etc.

(14) aliphatic dicarboxylic acids (JP-A Hei 1-56614; JP-A Hei 5-310565)

(15) silicides (JP-A Sho 62-249709)

25 (16) organogermanes (JP-A Hei 2-62885; JP-A Hei 5-255130; JP-A Hei 7-247296; JP-A Hei 8-59485)

(17) flavonoids (JP-A Hei 3-240725; JP-A Hei 7-206838; JP-A Hei 9-241165; WO 94/04520)

30 (18) alkylamines (JP-A Hei 6-206818; JP-A Hei 9-59233; JP-A Hei 9-40626; JP-A Hei 9-124471)

(19) amino acids (Published Japanese Translation of International Publication 4-502611; Published Japanese Translation of International Publication 7-503713)

35 (20) aromatic compounds such as ascochlorin (JP-A Hei 6-305959), benzoic acid (WO 91/11997), pyrrolonaphthyridinium (JP-A Hei 10-158265), etc.

(21) polypeptides (Published Japanese Translation of International Publication 7-500580)

(22) vitamins such as pyridoxamine (WO 97/09981), etc.

(23) SH group-containing compounds such as glutathione, cysteine,  
5 N-acetylcysteine, etc.

(24) SH group-containing proteins such as reduced albumin, etc.

(25) tetracyclines (JP-A Hei 6-256280)

(26) chitosans (JP-A Hei 9-221427)

(27) tannins (JP-A Hei 9-40519)

10 (28) quaternary ammonium ion-containing compounds

(29) biguanide agents such as phenformin, buformin, metformin,  
etc.

(30) ion exchange resins

15 (31) inorganic compounds such as activated charcoal, silica gel,  
alumina, calcium carbonate, etc.

20 The above compounds include those collectively known as Maillard  
reaction inhibitors. The Maillard reaction is a non-enzymatic  
glycation reaction between a reducing sugar such as glucose, and an  
amino acid or protein. Focusing on a phenomenon of brown coloration  
in a mixture consisting of amino acid and reducing sugar upon heating,  
Maillard reported this reaction in 1912 (Maillard, L. C., Compt. Rend.  
Soc. Biol., 72: 599 (1912)). This Maillard reaction is involved in  
brown coloration of food during heating or storage, generation of  
aromatic components and taste, and protein denaturation. Therefore,  
25 this reaction has been mainly studied in the field of food chemistry.

30 In 1968, glycated hemoglobin (HbA1c), a micro fraction of  
hemoglobin, was identified *in vivo*, which was revealed to increase  
in patients with diabetes (Rahbar. S., Clin. Chim. Acta, 22: 296 (1968)).  
These findings helped launch a wave of interest in the significance  
of *in vivo* Maillard reaction and the participation of the reaction  
in the onset of adult diseases such as diabetic complications and  
arteriosclerosis as well as the progress of aging. Agents inhibiting  
the *in vivo* Maillard reaction were explored intensively, resulting  
in the discovery of the above-mentioned compounds as agents inhibiting  
35 the Maillard reaction.

However, it was not known that such Maillard reaction inhibitors

are capable of improving carbonyl-stress state in peritoneal-dialysis patients and inhibit peritoneal dialysis-associated complications caused by carbonyl stress, by eliminating carbonyl compounds derived from the peritoneal dialysate and from the blood.

5       The inventive carbonyl compound-trapping agents include macro-molecular compounds such as ion exchange resins, or inorganic compounds such as activated charcoal, silica gel, alumina, and calcium carbonate, as well as organic compounds represented by the above-mentioned Maillard reaction inhibitors. These compounds are  
10       agents trapping insoluble carbonyl compounds in the peritoneal dialysate by utilizing their carbonyl compound-adsorbing activity. These compounds, though known to be chromatographic carriers, were not known to be useful for improving the carbonyl-stress state.

15       Adsorptive blood purifiers using activated charcoal have been in use for purifying blood in cases of drug poisoning or hepatic coma. They were also used in auxiliary treatment associated with hemodialysis for the removal of various endotoxins, exotoxins, and vasoactive substances that increase at the early stage of acute renal failure in multiple organ failure. However, it has been unknown that adsorptive  
20       blood purifiers are useful for removing carbonyl compounds present in the peritoneal dialysates or carbonyl compounds accumulated in the peritoneal cavity during the dialysis.

25       JP-A Sho 58-169458 describes an invention relating to a peritoneal dialysate containing a solid particulate absorbent and a method of peritoneal dialysis using this peritoneal dialysate. According to the publication, the solid particulate absorbent is added for the purpose of eliminating creatinine and low-molecular-weight metabolites; however, there is no description of effectiveness of the absorbent for the elimination of the carbonyl compounds accumulated  
30       in the peritoneal dialysate or in the peritoneal cavity during dialysis. Further, the publication does not indicate nor suggest that the carbonyl-stress state of peritoneal-dialysis patients can be improved by the method of peritoneal dialysis.

35       The composition of the peritoneal dialysate used as a base to which the inventive carbonyl compound-trapping agent is added, may be any conventional dialysate. A peritoneal dialysate generally

comprises an osmoregulatory agent, a buffering agent, and an inorganic salt. The sugars as listed above are used as osmoregulatory agents. In view of the stability of glucose, a buffer agent giving acidic pH (pH 5.0-5.4) is frequently used. Of course, when the osmoregulatory agent is not glucose, the buffering agent can be suitably selected to give a more physiological pH (pH of around 7.0). Alternatively, a product form has been designed, in which the buffering agent that adjusts pH at the time of use is packaged separately, allowing the use of both glucose and a neutral pH. In the present invention, carbonyl compounds generated in the processes of heat sterilization and long-term storage are eliminated, enabling the preferable use of a buffer system capable of giving neutral pH. Such a formulation has previously been difficult because of the degradation of glucose. A peritoneal dialysate generally contains inorganic salts such as sodium chloride, calcium chloride, or magnesium chloride. These salts bring peritoneal dialysates closer to physiological conditions, and are expected to give greater biocompatibility.

The inventive carbonyl compound-trapping agent can be added to a peritoneal dialysate of a known composition at the time of formulation, and the formulated and sealed dialysate can be sterilized by heating. It is expected that the addition of the agent sufficiently prevents the generation of carbonyl compounds from the major constituents during heat sterilization. Alternatively, the peritoneal dialysate is placed in a compartmentalized container comprising a first compartment and a second compartment; a reducing sugar is placed in the first compartment and the carbonyl compound-trapping agent is placed in the second compartment, and the two are mixed immediately before use. In this case, carbonyl compounds generated in the processes of sterilization and storage immediately bind to the carbonyl compound-trapping agent mixed. The excess carbonyl compound-trapping agents also trap blood-derived carbonyl compounds, after being administered into the peritoneal cavity. A single carbonyl compound-trapping agent or a combination of multiple trapping agents may be added to the peritoneal dialysate.

There can be many types of methods for contacting a peritoneal dialysate with a carrier on which a carbonyl compound-trapping agent

is immobilized, or with carbonyl compound-trapping agents that are insoluble in the peritoneal dialysate. For example, the peritoneal dialysate is enclosed in a container where the carbonyl compound-trapping agent is immobilized inside, or in a container carrying a carbonyl compound-trapping agent immobilized on particulates or fibrous carriers, thereby trapping carbonyl compounds that generate and accumulate during storage. In the latter system, the insoluble carriers can be separated from the peritoneal dialysate by filtration. Alternatively, a carbonyl compound-trapping cartridge is prepared by filling a column with carrier beads or fibrous carriers on which the carbonyl compound-trapping agent is immobilized, or with a carbonyl compound-trapping agent which per se is insoluble in the peritoneal dialysate. Then, the peritoneal dialysate is contacted with the carrier in the cartridge, and then the fluid is infused into the peritoneal cavity. It is preferred that distilled water is pre-filled in the cartridge to prevent air bubbles at the start of peritoneal dialysis. When the carbonyl compound-trapping cartridge is contacted with the peritoneal dialysate at the time of peritoneal infusion, although it is impossible to remove patient-derived carbonyl compounds that accumulate in the fluid during the dialysis, carbonyl compounds originally present in the dialysate can be eliminated. Alternatively, when peritoneal dialysis treatment is conducted by using a closed circuit where the peritoneal dialysate is circulated by a small circulating pump, it is possible to attain the removal of not only carbonyl compounds originally present in the dialysate but also those that accumulate in the peritoneal cavity during dialysis, by installing within the circuit, the above-mentioned carbonyl compound-trapping cartridge containing carriers with immobilized carbonyl compound-trapping agent.

There is no particular restriction on the carrier that is used for immobilizing the inventive carbonyl compound-trapping agent, as long as the carrier is harmless to the human body and is sufficiently safe and stable as a material that is directly in contact with the peritoneal dialysate. The carriers include, for example, synthetic or naturally-occurring organic macro-molecular compounds, and inorganic materials such as glass beads, silica gel, alumina, and

activated charcoal, as well as these materials coated with a polysaccharide or synthetic polymer. Conventional modifications, reformations or denaturations can improve the permeability, drug compatibility, protective strength, adsorption capacity, or carbonyl compound specificity of the carries on which the carbonyl compound-trapping agent is immobilized, or the carbonyl compound-trapping agent, which per se is insoluble in the peritoneal dialysate.

A carrier comprising a macromolecule is exemplified by a polymethyl methacrylate polymer, polyacrylonitrile polymer, polysulfone polymer, vinyl polymer, polyolefin polymer, fluorine polymer, polyester polymer, polyamide polymer, polyimide polymer, polyurethane polymer, polyacryl polymer, polystyrene polymer, polyketone polymer, silicon polymer, cellulose polymer, chitosan polymer; specifically, polysaccharides such as agarose, cellulose, chitin, chitosan, sepharose, dextran, etc. and derivatives thereof, and polyester, polyvinyl chloride, polystyrene, polysulfone, polyethersulfone, polypropylene, polyvinyl alcohol, polyarylether sulfone, polyacrylic ester, polymethacrylic ester, polycarbonate, acetylated cellulose, polyacrylonitrile, polyethylene terephthalate, polyamide, silicone resin, fluororesin, polyurethane, polyetherurethane, and polyacrylamide and derivatives thereof. The macromolecular material can be used alone or in a combination of two or more types of macromolecules. In the latter case, at least one of the macromolecules bears the carbonyl compound-trapping agent immobilized on it. The immobilized carbonyl compound-trapping agent is used alone or in a combination of two or more types of compounds. The above-mentioned polymer material may be a polymer comprising a single type of monomer or a copolymer comprising multiple types of monomers. Further, the material may be treated by the addition of an appropriate modifier, or may be subjected to denaturation treatment such as radiation cross-linking or cross-linking using peroxide.

There is no restriction on the shape of carrier. For example, the carrier can be membranous, fibrous, granular-shaped, hollow fiber-like, non-woven fabric-like, porous, or honeycomb-shaped. The carrier's area of contact with the peritoneal dialysate can be



controlled by varying the thickness, surface area, diameter, length, shape, and/or size of the carrier. Further, the carbonyl compound-trapping agent can be immobilized on the inner wall of container, where the peritoneal dialysate is placed, and also within the circuit where the peritoneal dialysate is circulated.

The carbonyl compound-trapping agent can be immobilized on the above-mentioned carrier by using known methods, such as physical adsorption, specific biochemical binding reaction, ion binding, covalent bonding, grafting, etc. If necessary, a spacer can be inserted between the carrier and the carbonyl compound-trapping agent. When the carbonyl compound-trapping agent exhibits toxicity, the amount released becomes a vital issue. Thus, it is preferred that the carbonyl compound-trapping agent is immobilized on the carrier by covalent bonding so as to minimize the released amount. Functional groups in the carrier are utilized for covalently bonding the carbonyl compound-trapping agent thereto. The functional group is, for example, hydroxyl group, amino group, aldehyde group, carboxyl group, thiol group, silanol group, amide group, epoxy group, succinylimino group, etc.; however, the functional group is not limited to these groups. The covalent bond is, for example, ester linkage, ether linkage, amino linkage, amid linkage, sulfide linkage, imino linkage, disulfide linkage or the like.

A commercially available product, for example polystyrene carrier having sulfonylhydrazine groups (PS-TsNHNH<sub>2</sub>, ARGONAUT TECHNOLOGIES CO.), can be used as a carrier for immobilizing carbonyl compound-trapping agent.

The carrier with the immobilized carbonyl compound-trapping agent of the present invention can be sterilized by an appropriate sterilization method selected from publicly known sterilization methods depending upon the types of carbonyl compound-trapping agent and carrier used. The sterilization method includes, for example, high-pressure steam sterilization, gamma-ray irradiation sterilization, gas sterilization, etc. A cartridge, which is filled with the insoluble carbonyl compound-trapping agent or carrier with immobilized carbonyl compound-trapping agent, is connected with a container containing the peritoneal dialysate to simultaneously

sterilize both the cartridge and container.

The carbonyl compound may be insufficiently eliminated from the peritoneal dialysate when only a small amount of carbonyl compound-trapping agent is in contact with the peritoneal dialysate.

5 In general, it is hard to predict the quantity of carbonyl compound present in the peritoneal dialysis. Accordingly, the carbonyl compound-trapping agent is used as much as possible without compromising the safety in patients so as to maintain the maximal effect. The dose of carbonyl compound-trapping agent can be controlled  
10 by altering the amount of carbonyl compound-trapping agent immobilized on the carrier or by altering the amount of carrier on which the carbonyl compound-trapping agent is immobilized at the time of use.

The carbonyl compound-trapping agent can also be infused into the peritoneal dialysis circuit through an appropriate mixing connector installed in the circuit. In this case, the carbonyl compound  
15 generated during sterilization and storage processes is trapped within the circuit.

Further, the carbonyl compound-trapping agent can be directly administered into the peritoneal cavity and mixed with the peritoneal dialysate in the peritoneal cavity. In this case, carbonyl compounds  
20 derived from the peritoneal dialysate and from the blood are inactivated in the peritoneal cavity.

Furthermore, prior to the infusion of peritoneal dialysate to a patient, or while the fluid is dwelling in the peritoneal cavity,  
25 the carbonyl compound-trapping agent is administered to the peritoneal-dialysis patient by intravenous injection or the like, thereby successfully achieving the improvement of carbonyl-stress state in the peritoneal cavity.

Preferred embodiments of the inventive peritoneal dialysate are  
30 specifically described below.

The composition of the base dialysate is generally as follows.

Glucose	1-5%w/v
Sodium ion	100-150 meq
Potassium ion	0-0.05 meq
Magnesium ion	0-2 meq
Calcium ion	0-4 meq

Chloride ion 80-150 meq

Buffering agent 10-50 mM

(organic acids such as lactic acid, citric acid, malic acid, acetic acid, pyruvic acid, and succinic acid)

- 5 This is only a general formula, and a more suitable composition is selected depending on the symptoms of the patient.

10 The inventive carbonyl compound-trapping agent is added in an effective amount to the above-indicated basic formula. For example, when aminoguanidine is used, the concentration is 1 mM or higher, preferably 10 mM or higher, more preferably 10 mM or higher but not higher than 100 mM. If the amount of carbonyl compound-trapping agent added is small, they might be used up for the carbonyl compounds generated in the processes of production and storage. As a consequence, the trapping agent is unable to treat carbonyl compounds transferred to the dialysate from the blood and tissues of a patient during dialysis. Particularly, it is hard to predict the quantity of carbonyl compounds transferred from the blood and tissues to the dialysate. Accordingly, the carbonyl compound-trapping agent is used as much as possible without compromising the safety of a patient so as to maintain the maximal effect. It has been known that aminoguanidine exhibits only a low toxicity to animals. According to "Registry of Toxic Effect of Chemical Substances" (1978), the half-lethal dose of aminoguanidine, which is subcutaneously administered, is 1258 mg/kg in rat, or 963 mg/kg in mouse. This compound is water-soluble. OPB-9195 can also be added similarly, to a concentration of 1 mM or higher, preferably 10 mM or higher, more preferably 10 mM or higher, but not higher than 100 mM.

20 The inventive peritoneal dialysate as formulated above is filled in an appropriate closed container, and subjected to sterilization. 30 An effective sterilization includes heat sterilization such as high-pressure steam sterilization, and hot water sterilization. In this case, it is important to use a container that releases no toxic substances at a high temperature, and has enough strength to bear transportation after sterilization. A specific example of such a 35 container is a flexible plastic bag made of polyvinyl chloride, polypropylene, polyethylene, polyester, or ethylene-vinyl acetate

copolymer. Further, to avoid the deterioration of the fluid resulting from the influence of the outside air, the container filled with the peritoneal dialysate may be packaged using a high gas barrier packaging material.

5 Filter sterilization can be selected in place of heat sterilization. For example, the fluid is sterilized by filtration using a precision filter device with a membrane of about 0.2- $\mu$ m pore size. This method is free of the generation of carbonyl compounds that result during heating. The filter-sterilized peritoneal  
10 dialysate is filled into a container such as a flexible plastic bag and then sealed. Because the series of processes from sterilization to transportation does not differ from the preparation of current dialysates, the inventive peritoneal dialysate can also be manufactured by a procedure comprising similar steps.

15 When the sterilization is achieved by heat sterilization including high-pressure heat sterilization, as long as the carbonyl compound-trapping agent used is sufficiently stable to a treatment such as heating, the trapping agent may be added when the peritoneal dialysate is formulated, prior to the heat sterilization process.  
20 This procedure reduces the generation and accumulation of dialysate-derived carbonyl compounds during heating. Of course, the carbonyl compound-trapping agent also functions to reduce the generation and accumulation of carbonyl compounds during storage and peritoneal dialysis.

25 When the carbonyl compound-trapping agent used is unstable to heat sterilization, it can be sterilized by a method that does not require heating. Such sterilization methods include, for example, filter sterilization. Alternatively, the carbonyl compound-trapping agent may be added to a sterilized peritoneal dialysate. There is  
30 no particular limitation on the timing of addition. For example, the carbonyl compound-trapping agent is preferably added to the fluid after sterilization, because in this case, the trapping agent can suppress not only the generation of carbonyl compounds during peritoneal dialysis, but also the generation and accumulation of  
35 carbonyl compounds in the peritoneal dialysate during the storage prior to dialysis.

Alternatively, the carbonyl compound-trapping agent can be added immediately before peritoneal dialysis treatment or at the time of treatment. For example, the base solution and the carbonyl compound-trapping agent are placed separately in the above-mentioned flexible plastic bags and such, before the treatment, and then, the two are mixed together in sterile conditions at the start of peritoneal dialysis. To achieve this, a flexible plastic bag as disclosed in Unexamined Published Japanese Patent Application No. (JP-A) Sho 63-19149, which comprises two compartments separated by a removable partition, is suitably used.

Alternatively, a connector for mixing is installed in the peritoneal dialysis circuit, and the carbonyl compound-trapping agent can be infused through the connector.

This type of preparation procedure for peritoneal dialysates can be used to prepare heat-stable carbonyl compound-trapping agents as well as heat-unstable carbonyl compound-trapping agents.

Further, when peritoneal dialysis is conducted by circulating the peritoneal dialysate in a closed circuit with a small circulating pump, a filter device filled with the carbonyl compound-trapping agent may be installed anywhere in the circuit.

The peritoneal dialysate of the present invention can be used for a peritoneal dialysis treatment similar to a treatment using currently used peritoneal dialysates. Specifically, a suitable amount of the inventive peritoneal dialysate is infused into the peritoneal cavity of a patient, and low-molecular weight constituents present in the body are allowed to transfer into the peritoneal dialysate through the peritoneum. The peritoneal dialysate is circulated intermittently, and the treatment is further continued depending on the symptoms of the patient. During this period, the carbonyl compounds, as well as other substances such as creatinine, inorganic salt, and chloride ion, transfer to the peritoneal dialysate from blood and from inside the peritoneum. At the same time, the toxic activity of carbonyl compounds is eliminated by the carbonyl compound-trapping agent, thereby rendering the compounds harmless.

### Brief Description of the Drawings

Figure 1 shows the amounts of carbonyl compounds present in peritoneal dialysates and the effluent fluids.

Figure 2 shows a photograph of histological localization of carbonyl-modified proteins in the peritoneal tissue of a peritoneal-dialysis patient (in the top panel) and the corresponding schematic illustration (in the bottom panel). In this figure, A indicates positive areas in the connective tissues, on which mesothelial cells had been present but have been removed; B indicates positive areas on the thickened vascular wall.

Figure 3 shows VEGF mRNA expression in mesothelial cells exposed to glyoxal, methylglyoxal, and 3-deoxyglucosone. Reverse transcription was performed on total RNA from cultured rat mesothelial cells incubated with various concentrations (0, 100, 200, and 400  $\mu$ M) of glyoxal (A), 3-deoxyglucosone (B), or methylglyoxal (C). VEGF and G3PDH cDNAs were amplified by PCR for 30 and 21 cycles, respectively. Experiments were performed in triplicate to calculate averages. Ratio of VEGF mRNA to G3PDH mRNA was calculated for the average of each experiment. The average  $\pm$  S.D. of the three experiments is illustrated in the figure. \* $P < 0.0005$ .

Figure 4 shows VEGF protein production in microvascular endothelial cells exposed to methylglyoxal. Human endothelial cells were cultured in the presence of various concentrations (0, 200, and 400  $\mu$ M) of methylglyoxal, and VEGF protein released into culture supernatant was quantified by ELISA. Representative data from the three experiments are shown. The data are expressed as mean  $\pm$  range. \* $P < 0.05$ , and \*\* $P < 0.01$ .

Figure 5 shows VEGF mRNA expression in endothelial cells exposed to methylglyoxal. Human endothelial cells were cultured in the presence of various concentrations (0, 100, 200, and 400  $\mu$ M) of methylglyoxal. VEGF and G3PDH cDNAs were amplified by PCR for 30 and 21 cycles, respectively. Experiments were performed in triplicate to calculate mean values. Ratio of VEGF mRNA to G3PDH mRNA was calculated for the mean of each experiment. The mean  $\pm$  S.D. of the three experiments is illustrated in the figure. \* $P < 0.05$ , \*\* $P < 0.005$ , and \*\*\* $P < 0.0001$ .

Figure 6 shows VEGF mRNA expression in peritoneal tissues of rats given daily intraperitoneal loads of methylglyoxal for 10 days. VEGF and G3PDH mRNA expressed in peritoneums were amplified by RT-PCR for 28 and 16 cycles, respectively. Experiments were performed in triplicate to calculate mean values. Ratio of VEGF mRNA to G3PDH mRNA was calculated for the mean of each experiment. The mean  $\pm$  S.D. of the three experiments is illustrated in the figure. \*P<0.05.

Figure 7 shows the generation process of carbonyl stress in the peritoneal cavity of peritoneal-dialysis patient.

Figure 8 shows the effect of the addition of aminoguanidine on the generation of pentosidine following the incubation of the peritoneal dialysis effluent.

Figure 9 shows the effect of the addition of aminoguanidine on the generation of protein carbonyl following the incubation of the peritoneal dialysis effluent.

Figure 10 shows the effect of the addition of aminoguanidine on the amounts of carbonyl compounds present in peritoneal-dialysis (CAPD) effluents from three peritoneal-dialysis patients (patient I, patient S, and patient K).

Figure 11 shows the effect of the addition of aminoguanidine on the generation of 5-HMF within the peritoneal dialysate in an acidic pH range.

Figure 12 shows the effect of the addition of aminoguanidine on the generation of 5-HMF within the peritoneal dialysate in a neutral pH range.

Figure 13 shows the effect of the addition of aminoguanidine on the amounts of carbonyl compounds within the peritoneal dialysate in acidic and neutral pH ranges.

Figure 14 shows the suppressing effect of carbonyl compound-trapping beads added to the peritoneal dialysate on the generation of pentosidine.

Figure 15 shows the carbonyl compound-eliminating effect of carbonyl compound-trapping beads added to the peritoneal dialysate.

Figure 16 shows the carbonyl compound-eliminating effect of activated charcoal added to a dicarbonyl compound solution.

Figure 17 shows the carbonyl compound-eliminating effect of

activated charcoal added to the peritoneal dialysate.

Figure 18 shows the trapping activity of guanidine for glyoxal, methylglyoxal, and 3-deoxyglucosone.

Figure 19 shows the trapping activity of metformin, which is a biguanide, for glyoxal, methylglyoxal, and 3-deoxyglucosone.

Figure 20 shows the trapping activity of buformin, which is a biguanide, for glyoxal, methylglyoxal, and 3-deoxyglucosone.

Figure 21 shows the trapping activity of phenformin, which is a biguanide, for glyoxal, methylglyoxal, and 3-deoxyglucosone.

Figure 22 shows the trapping activity of aminoguanidine for glyoxal, methylglyoxal, and 3-deoxyglucosone.

Figure 23 shows the trapping activity of cysteine, which is an SH compound, for glyoxal, methylglyoxal, and 3-deoxyglucosone.

Figure 24 shows the trapping activity of N-acetylcysteine, which is an SH compound, for glyoxal, methylglyoxal, and 3-deoxyglucosone.

Figure 25 shows the trapping activity of GSH, which is an SH compound, for glyoxal, methylglyoxal, and 3-deoxyglucosone.

Figure 26 shows the trapping activity of albumin, which is an SH compound, for glyoxal, methylglyoxal, and 3-deoxyglucosone.

Figure 27 shows the suppressing effect of SH compounds added to the peritoneal dialysate on the generation of pentosidine.

#### Best Mode for Carrying out the Invention

The present invention is illustrated more specifically below with reference to examples, but is not to be construed as being limited thereto.

#### Example 1. Measurement of the amount of carbonyl compounds present in the peritoneal dialysate and peritoneal dialysate effluent

In order to demonstrate the generation of carbonyl stress in the peritoneal cavity, the amount of carbonyl compounds present in the peritoneal dialysate effluent was measured according to the following experiment method.

##### (i) Measurement of carbonyl compounds

After the peritoneal dialysate (Baxter Ltd.; Dianeal PD-2 2.5) had been administered to a peritoneal-dialysis patient and had been



allowed to dwell in the peritoneal cavity overnight, the peritoneal dialysate effluent was collected from the patient. Aliquots (400  $\mu$ l) of both the peritoneal dialysate and the peritoneal dialysate effluent were separately mixed with a 400- $\mu$ l solution of 0.5 N hydrochloric acid containing 1.5 mM 2,4-dinitrophenylhydrazine (2,4-DNPH) (Wako Pure Chemical Industries, Ltd.), and the mixture was stirred at room temperature for 30 minutes to react the carbonyl compound with 2,4-DNPH. Subsequently, an aqueous solution of 1 M acetone (40  $\mu$ l) was added to the mixture, which was then stirred at room temperature for 5 minutes to remove excess 2,4-DNPH. The mixture was washed three times with 400  $\mu$ l of n-hexane. The aqueous layer was recovered and the absorbance thereof was measured at 360 nm in a spectrophotometric microplate reader (Nippon Molecular Devices Co.; SPECTRAMax250).

(ii) Preparation of calibration curve

Aqueous solutions of various glucose concentrations were prepared and the amounts of carbonyl compounds derived from glucose were measured by the same method as described in (i). A calibration curve of glucose concentration vs. concentration of carbonyl compound was prepared based on this experiment.

(iii) Quantification of carbonyl compounds

Respective glucose concentrations of the peritoneal dialysate and peritoneal dialysate effluent were measured by using a glucose assay kit (Wako Pure Chemical Industries, Ltd.; Glucose CII-Test Wako). The amount of carbonyl compounds derived from glucose was estimated by using the calibration curve. The amount of carbonyl compounds in the fluids was determined by subtracting the amount of glucose-derived carbonyl compounds from the total amount of carbonyl compound in the sample solutions.

The result obtained is shown in Fig. 1. The peritoneal dialysate effluent, which had dwelled overnight in the peritoneal cavity, contained five times more carbonyl compounds than the peritoneal dialysate prior to the administration did. This indicates the transfer of carbonyl compounds from the blood into the peritoneal cavity.

Example 2. Histological localization of carbonyl-modified proteins in the peritoneum of a peritoneal-dialysis patient

The localization of carbonyl compounds in peritoneal tissues of a peritoneal-dialysis patient was studied by immunostaining using malondialdehyde as the index.

Peritoneal tissues from a peritoneal-dialysis patient (50-year-old male with a five-year history of peritoneal dialysis) were processed by immunostaining according to the method of Horie et al. (Horie, K. et al., J. Clin. Invest., 100, 2995-3004 (1997)). The primary antibody used was a mouse anti-malondialdehyde monoclonal antibody. The results showed that intense positive signals were present in the connective tissue under removed mesothelial cells, and the signals were also present on the thickened vascular wall (Fig. 2).

Malondialdehyde (MDA) is a carbonyl compound, which is generated by the degradation of peroxide lipid. Therefore, immunostaining was carried out by using antibodies against 4-hydroxy-2-nonenal, a degradation product of peroxide lipid other than MDA, and against carboxymethyllysine and pentosidine, which are generated by the oxidation of saccharides. The results showed that the positive signals were localized in the same areas as those shown in Fig. 2. These indicate that, in the peritoneal tissues of peritoneal-dialysis patients, proteins are modified in association with the enhancement of carbonyl stress caused by the presence of degradation products of peroxide lipids and carbonyl compounds generated by the oxidation of saccharides.

Example 3. Peritoneal cell damages by methylglyoxal

Decreased ultrafiltration of peritoneum by long-term peritoneal dialysis has been taken as evidence for an augmentation of the peritoneal surface area available for diffusive exchange (Krediet RT, Kidney Int, 55: 341-356 (1999); Heimbürger O et al., Kidney Int, 38: 495-506 (1990); Imholz AL et al., Kidney Int, 43: 1339-1346 (1993); Ho-dac-Pannekeet M et al., Perit Dial Int, 17: 144-150 (1997)). Namely, glucose in peritoneal dialysate flows out of peritoneal cavity by diffusion, thereby decreasing the dialytic function according to

osmotic pressure gradient. An increased vascular surface area within the peritoneum might also account for this. In addition, the vascular endothelial growth factor (VEGF) might play a critical role in this pathology. VEGF increases vascular permeability (Senger DR et al., Science 219: 983-985 (1983); Connolly DT et al., J Biol Chem, 264: 20017-20024 (1989)), stimulates nitric oxide (NO) synthesis and vasodilation (Hood JD et al., Am J Physiol, 274: H1054-1058 (1998)), and induces inflammatory responses (Clauss M et al., J Exp Med, 172: 1535-1545 (1990); Melder RJ et al., Nat Med, 2: 992-997 (1996)). Furthermore, VEGF is a powerful angiogenic factor contributing to the recovery from vascular lesions (Thomas KA, J Biol Chem, 271: 603-606 (1996); Ferrara N et al., Endocr Rev, 18: 4-25 (1997); Shoji M et al., Am J Pathol, 152: 399-411 (1998)). Consequently, effects of glucose degradation products contained in dialysate on VEGF production were examined.

<3-1> VEGF expression in peritoneum mesothelial and vascular endothelial cells cultured in the presence of glucose degradation products

Peritoneum was obtained from 6-week-old male CD (SD) IGS rats (Charles-River, Kanagawa, Japan). Mesothelial cells were isolated based on the method of Hjelle et al. (Hjelle JT et al., Perit Dial Int, 9: 341-347 (1989)) and cultured in Dulbecco's modified Eagle's medium (DMEM) containing 10% fetal bovine serum. The mesothelial cells at passage 7 to 10 were cultured for 3 hours in a CO<sub>2</sub> incubator in the presence of various concentrations of glucose degradation products (glyoxal, methylglyoxal (Sigma, St. Louis, MO), or 3-deoxyglucosone (kindly provided from Fuji Memorial Research Institute, Otsuka Pharmaceutical Co., Kyoto, Japan)). The expression of VEGF mRNA was analyzed by semiquantitative RT-PCR. Total RNA was isolated from the mesothelial cells using Rneasy Mini Kit (Qiagen, Germany). Five micrograms of the RNA was reverse-transcribed using oligo(dT)<sub>12-18</sub> primers (Gibco BRL, Gaithersburg, MD) with 200 units of RNase H-free reverse transcriptase (Superscript II: Gibco BRL) and PCR amplification was performed as described previously (Miyata T et al., J Clin Invest, 93: 521-528 (1994)). The sequences of the primers used for the amplification of rat VEGF were 5'-ACTGGACCCTGGCTTTACTGC-3'

(SEQ ID NO: 1) and 5'-TTGGTGAGGTTTGATCCGCATG-3' (SEQ ID NO: 2). The amplified product was 310 bp long. The primers used for the amplification of rat glyceraldehyde-3-phosphate dehydrogenase (G3PDH) were 5'-CCTGCACCACCAACTGCTTAGCCC-3' (SEQ ID NO: 3) and 5'-GATGTCATCATATTTGGCAGGTT-3' (SEQ ID NO: 4) and amplified a 322 bp fragment. The G3PDH served as an internal RNA control to allow comparison of RNA levels among different specimens. Specimens were amplified in a DNA Thermal cycler (Perkin Elmer Cetus, Norwalk, CT) to determine the suitable number of cycles consisting of 0.5 min at 94°C, 1 min at 60°C, and 1.5 min at 72°C. In preliminary experiments, reverse transcription and PCR amplification were performed on various amounts of RNA for 16, 18, 21, 25, 28, 31, and 34 cycles. These experiments showed that, with 30 cycles of amplification for VEGF mRNA and with 21 cycle of G3PDH mRNA amplification, PCR product signals were quantitatively related to input RNA. PCR products resolved by electrophoresis in 1.5% agarose gel and stained with ethidium bromide were quantified by measuring the signal intensity with a quantification program (NIH image). Experiments were performed for each glucose degradation product concentration. Messenger RNA was determined in triplicate and the results were averaged for each experiment. A total of 3 to 4 independent experiments were performed for each experimental condition. The results were averaged and expressed as mean  $\pm$  S.D. The statistical significance was evaluated by analysis of variance (ANOVA). If a significant difference was indicated by this analysis, results obtained with different concentrations of methylglyoxal were compared by Scheffe's t-test.

At concentrations varying from 0 to 400  $\mu$ M, neither glyoxal nor 3-deoxyglucosone modified VEGF expression (Figs. 3A and 3B). Only methylglyoxal stimulated VEGF mRNA expression at a concentration of 400  $\mu$ M ( $P < 0.0005$ ) (Fig. 3C). The sample RNAs that had not been reverse-transcribed did not yield the PCR product. All cells remained viable. Mesothelial cells were also cultured in the presence of higher concentrations of 3-deoxyglucosone (0.625, 2.5, and 5 mM), showing a decreased viability (cell viability was 80, 55, and 8% in the presence of 0.625, 2.5, and 5 mM 3-deoxyglucosone, respectively). Because of the decreased viability, VEGF mRNA expression could not be measured.

As a consequence of these observations, only methylglyoxal was used in the subsequent experiments.

The release of VEGF protein into the culture supernatant was measured by ELISA for mesothelial cells cultured for 24 hours in the presence of various concentrations of methylglyoxal. Human microvascular endothelial cells were purchased from Kurabo (Osaka, Japan) and cultured in VEGF-depleted EGM-2 medium (Takara, Tokyo, Japan). The cells were incubated with methylglyoxal in the same manner as rat peritoneum mesothelial cells. The VEGF protein in the culture supernatant prepared in duplicate was quantified by enzyme-linked immunosorbent assay (ELISA) using a kit (Quantikine: R&D Systems, Minneapolis, USA) according to the attached manual. The experiment was repeated three times. The results were statistically analyzed as described above.

As a result, addition of methylglyoxal to the medium resulted in a dose-dependent increase of VEGF (Fig. 4). No VEGF release was detected during in the absence of methylglyoxal. Namely, at protein level, methylglyoxal also dose-dependently stimulates the production and release of VEGF.

VEGF mRNA expression was then assessed in endothelial cells cultured in the presence of various concentrations (0 to 400  $\mu$ M) of methylglyoxal. Experiments were performed as those for rat mesothelial cells. However, primers used for human VEGF amplification were 5'-GGCAGAATCATCACGAAGTGGTG-3' (SEQ ID NO: 5) and 5'-CTGTAGGAAGCTCATCTCTCC-3' (SEQ ID NO: 6). The amplified fragment was 271 bp long. The same primers were used for human and rat G3PDH amplifications. The analyses revealed that VEGF mRNA expression rose in a dose-dependent manner (Fig. 5).

<3-2> VEGF expression in peritoneal tissues of rats given intraperitoneal injection of methylglyoxal

To further assess the biological effects of methylglyoxal on VEGF mRNA expression in the peritoneum *in vivo*, rats were given various amounts of methylglyoxal into their peritoneal cavity for 10 days. Six-week-old male CD (SD) IGS rats were given a daily intraperitoneal injection of 50 ml/kg of a saline solution containing various concentrations of methylglyoxal for 10 days. Peritoneum was isolated

from the parietal walls and investigated for VEGF mRNA expression. Experiments were performed as the above-mentioned *in vitro* experiments for rat mesothelial cells. However, mRNA was extracted from peritoneal tissues with ISOGEN (Nippon Gene, Tokyo, Japan). In addition, PCR amplification was performed for 28 cycles for VEGF mRNA and for 16 cycles for G3PDH mRNA.

As shown in Fig. 6, VEGF mRNA expression in samples of the parietal peritoneum increased significantly ( $P < 0.05$ ) according to methylglyoxal concentration. On optical microscopy, the peritoneal tissue was unaffected: the number of vessels, the vascular wall, the interstitium, and mesothelial cells remained normal.

<3-3> VEGF and carboxymethyllysine (CML) immunostaining of peritoneal tissue of long-term peritoneal dialysis patients

The distribution of VEGF and carboxymethyllysine was examined by immunohistochemistry in the peritoneal tissues of nine peritoneal dialysis patients. Carboxymethyllysine is derived from glucose degradation products such as glyoxal and 3-deoxyglucosone (Miyata T et al., *Kidney Int*, 55: 389-399 (1999)). An anti-carboxymethyllysine antibody was therefore used as a marker of glucose degradation product-modified proteins.

Peritoneal tissues were isolated, after obtaining informed consent, from nine non-diabetic peritoneal dialysis patients during catheter reinsertion (Table 1). Reinsertion was necessitated by catheter failure due to damage, incorrect position, and/or obstruction. No patients suffered from peritonitis. Normal peritoneal tissue was obtained, during abdominal surgery, from two male subjects (48 and 58 years old) with normal renal function.

Two- $\mu$ -thick peritoneal tissue sections were mounted on slides coated with 3-aminopropyltriethoxy silane (Sigma), deparaffined, rehydrated in distilled water, and incubated with a buffer solution (0.05 M Tris-HCl (pH 7.2), 0.1 M NaCl) containing Pronase (0.5 mg/ $\mu$ l: Dako, Glostrup, Denmark) for 15 min at room temperature. The slides were washed with PBS containing 0.5% Tween 20, blocked in 4% skim milk for 2 hours, and subsequently incubated with anti-VEGF rabbit IgG (Santa Cruz Biotechnology, Santa Cruz, CA) or anti-AGE mouse IgG (Ikeda K et al., *Biochemistry*, 35: 8075-8083 (1996)), the epitope

of which was carboxymethyllysine (Miyata T et al., Kidney Int, 51: 1170-1181 (1997)), overnight in humid chambers at 4°C. The sections were washed and incubated with 1:100 diluted goat anti-rabbit IgG conjugated with peroxidase or goat anti-mouse IgG conjugated with peroxidase (Dako) for 2 hours at room temperature, followed by the detection with 3,3'-diaminobenzidine solution containing 0.003% H<sub>2</sub>O<sub>2</sub>. Periodic acid-Schiff staining was also performed for histological analysis. Immunostaining was independently evaluated for signal intensity and distribution by two observers.

Results are shown in Table 1. In the table, "normal1" and "normal2" indicate samples derived from the subjects with normal renal function and "PD1" to "PD9" indicate samples derived from peritoneal dialysis patients. In the pictures of the peritoneal tissue of a representative long-term peritoneal dialysis patient (PD6 in Table 1), interstitial fibrosis, and thickening and hyalinosis of vascular walls were observed. Both VEGF and carboxymethyllysine co-localized in the mesothelial cells and vascular walls. In mesothelial layer, the signal with VEGF was weaker than that with carboxymethyllysine. Results were similar for the eight other patients. In the normal peritoneal sample (normal2 in Table 1), in contrast with peritoneal dialysis samples, VEGF was present only in the vascular walls but was absent in the mesothelial layer. Carboxymethyllysine was absent in the mesothelial layer and the signal was very weak in the vascular walls. Observations were similar in another control normal sample. No immunostaining was observed when normal mouse IgG was used. Thus, the fact that VEGF expression in the mesothelial layers and its co-localization with carboxymethyllysine are observed only in peritoneal dialysis patients suggests that the glucose degradation products present in peritoneal dialysate actually enhance VEGF production in uremic patients through peritoneal dialysis.

Table 1 . Immunohistochemical detection of CML and VEGF in peritoneal tissues of PD patients

Samples	Gender	Age (yr)	PD duration (months)	CML		VEGF	
				mesothelial layer	vascular walls	mesothelial layer	vascular walls
normal1	M	48	-	-	±	-	+
normal2	M	58	-	-	±	-	+
PD1	F	53	3	+	+	+	+
PD2	M	44	4	+	+	+	+
PD3	M	43	45	+	+	+	+
PD4	F	54	60	++	++	++	+
PD5	M	52	70	++	++	+	+
PD6	M	51	90	++	++	++	++
PD7	M	45	105	++	++	++	++
PD8	M	62	108	++	++	++	++
PD9	M	66	110	++	++	++	++

-: negative, ±: faint, +: positive, ++: strongly positive

The results mentioned above confirm that peritoneal dialysis patients are in carbonyl-stress state because of glucose degradation products or the like in peritoneal dialysate and further demonstrate for the first time that methylglyoxal enhances VEGF production in peritoneal cells. This suggests that at least a part of causes for decrease of peritoneal permeability by glucose degradation products contained in peritoneal dialysate is enhancement of VEGF production and angiogenic stimulation accompanied with it.

#### Example 4. Effect of a carbonyl compound-trapping agent added to dialysate effluents from peritoneal-dialysis patients

<4-1>

Pentosidine is an AGE structure, which has been seen to accumulate 20 times more in the blood of patients with renal failure compared to normal healthy persons (Miyata, T. et al., J. Am. Soc. Nephrol., 7: 1198-1206 (1996)). The inventors tested how aminoguanidine addition influenced the increase of pentosidine and carbonyl group on proteins (protein carbonyl) when the fluid effluent from a peritoneal-dialysis patient was incubated at 37°C.

An effluent from a peritoneal-dialysis patient, which had been kept overnight, was centrifuged, and the resulting supernatant was



sterilized by filtration (pore size; 0.45  $\mu$ m). Aminoguanidine (Tokyo Kasei Kogyo Co.) was added to the fluid at a final concentration of 0, 1, 10, or 100 mM. The mixture was incubated at 37°C. The incubation period was 1 to 2 when assaying pentosidine, and 2 weeks when assaying protein carbonyl. The quantitative assay for pentosidine was performed using HPLC (SHIMADZU Co.; LC-10A) after the hydrolysis of proteins in 6 N HCl at 110°C (T. Miyata et al., 1996, J. Am. Soc. Nephrol, 7:1198-1206; T. Miyata et al., 1996, Proc. Natl. Acad. Sci. USA, 93:2353-2358). The quantitative determination of protein carbonyl was carried out by measuring the absorbance (Nippon Molecular Devices Co.; SPECTRAMax 250) of hydrazone generated by the reaction of a carbonyl group after incubation with 2,4-dinitrophenylhydrazine (2,4-DNPH; Wako Pure Chemical Industries, Ltd.) (Levine, R. L. et al., Methods Enzymol., 233, 346-357 (1994)).

Based on the experimental result, aminoguanidine was shown to inhibit the generation of pentosidine in a concentration-dependent manner (Fig. 8). Aminoguanidine exhibited a similar inhibiting effect on the generation of protein carbonyl in a concentration-dependent manner (Fig. 9).

<4-2>

Next, aminoguanidine was tested for its effect on the amounts of carbonyl compounds other than glucose present in a fluid that was kept overnight in a peritoneal-dialysis patient, according to the following experiment method.

#### (A) Incubation of the peritoneal dialysate effluent

The peritoneal dialysate effluent was recovered from the patient and filtered with a filter with a pore size of 0.45  $\mu$ m. Aminoguanidine (Tokyo Kasei Kogyo Co.) was added to the fluid at a concentration of 0, 10, 50, or 250 mM to prepare a sample solution. An aliquot (1 ml) of the sample solution was added into a plastic tube with a screw cap, and the tube was incubated at 37°C for 15 hours. The sample solutions were stored at -30°C prior to the incubation.

#### (B) Quantification of carbonyl compounds

##### (i) Assay for carbonyl compounds present in sample solutions

Each of 400- $\mu$ l aliquots of sample solutions was mixed with a 400- $\mu$ l solution of 0.5 N hydrochloric acid containing 1.5 mM 2,4-DNPH (Wako

Pure Chemical Industries, Ltd.), and then, the mixture was stirred at room temperature for 30 minutes to react the carbonyl compound with 2,4-DNPH. Subsequently, an aqueous solution of 1 M acetone (40  $\mu$ l) was added to the mixture. The resulting mixture was stirred at room temperature for 5 minutes to remove excess 2,4-DNPH by reacting it with acetone. The aqueous mixture was washed three times with 400  $\mu$ l of n-hexane. The aqueous layer was recovered and the absorbance thereof was measured at 360 nm in a spectrophotometric microplate reader (Nippon Molecular Devices Co.; SPECTRAMax250).

(ii) Preparation of calibration curve

Aqueous solutions of various glucose concentrations were prepared and the amounts of carbonyl compounds derived from glucose were measured by the same method as described in (i); a calibration curve of glucose concentration vs. concentration of carbonyl compound was prepared based on this experiment.

(iii) Quantification of carbonyl compounds

Respective glucose concentrations of the samples were determined by using a glucose assay kit (Wako Pure Chemical Industries, Ltd.; Glucose CII-Test Wako). The amount of carbonyl compounds derived from glucose was estimated by using the calibration curve. The amount of carbonyl compound in the sample was determined by subtracting the amount of glucose-derived carbonyl compounds from the total amount of carbonyl compounds in the sample solution.

The result obtained is shown in Fig. 10. As the concentration of aminoguanidine increased, the amounts of carbonyl compounds other than glucose decreased, both in the unincubated and incubated (37°C for 15 hours) samples.

These results showed that the addition of the carbonyl compound-trapping agent to the peritoneal dialysate or the administration of the trapping agent to patients is effective in inhibiting the generation and/or accumulation of carbonyl compounds in the peritoneal dialysate infused into the peritoneal cavity. Thus, carbonyl compounds derived from the peritoneal dialysate or from the blood are eliminated from the peritoneal cavity, thereby achieving the improvement of the carbonyl-stress state in peritoneal-dialysis patients.

Example 5. Effect of a carbonyl compound-trapping agent added to the peritoneal dialysate in the process of heat sterilization

Peritoneal dialysates contain a high concentration of glucose as an osmoregulatory agent (1.35-4.0 w/v%). Glucose is unstable to heating and undergoes degradation during heat sterilization or storage. Degradation products of glucose have been reported to include 5-hydroxymethylfurfural (5-HMF), levulinic acid, and acetaldehyde (Richard, J. U. et al., Fund. Appl. Toxic., 4: 843-853(1984), Nilsson, C. B. et al., Perit. Dial. Int., 13: 208-213(1993)). The inventors tested aminoguanidine for its inhibiting effect on the generation of carbonyl compounds during heat sterilization of a peritoneal dialysate, by monitoring the quantities of 5-HMF and carbonyl compounds. Since the degree of glucose degradation is affected by pH of the solution, two peritoneal dialysates of acidic pH (pH 5.3) and neutral pH (pH 7.0) were prepared to carry out the experiment. Sterilization temperature was 121°C.

<5-1>

5-HMF was assayed by high performance liquid chromatography (SHIMADZU Co.; LC-10A) (Nilsson, C. B. et al., Perit. Dial. Int., 13: 208-213(1993)).

The results showed that aminoguanidine effectively inhibited the generation of 5-HMF in a concentration-dependent manner at both acidic pH (Fig. 11) and neutral pH (Fig. 12).

<5-2>

Quantification of carbonyl compounds in the peritoneal dialysate was carried out in the same manner as described in Example 1 by measuring the absorbance after the reaction with 2,4-DNPH (Levine, R. L. et al., Methods Enzymol., 233: 346-357 (1994)). Sterilization temperature was 121°C.

The result showed that aminoguanidine also effectively inhibited the generation of carbonyl compounds in a concentration-dependent manner (Fig. 13).

These findings clarified that the addition of an agent inhibiting the generation of carbonyl compounds to the peritoneal dialysate is highly effective in inhibiting the generation and/or accumulation

of carbonyl compounds in the peritoneal dialysate.

Example 6. Effect of carbonyl compound-trapping beads added to the peritoneal dialysate on the generation of pentosidine

5 Carbonyl compound-trapping beads, which comprise  
sulfonylhydrazine group-bound crosslinked polystyrene resin  
(PS-TsNHNH<sub>2</sub>; ARGONAUT TECHNOLOGIES CO.), were tested for the effect  
of eliminating carbonyl compounds from the peritoneal dialysate. The  
peritoneal dialysate, and the dialysate containing carbonyl  
10 compound-trapping beads, were incubated at 37°C to evaluate the effect  
of inhibiting the generation of pentosidine. Dimethylsulfoxide (100  
μl) was added to the tube containing carbonyl compound-trapping beads,  
to swell the beads. Then 800 μl of a peritoneal dialysate (Baxter  
Ltd.; Dianeal PD-4,1.5) and 200 μl of an aqueous solution of 150 mg/ml  
15 bovine serum albumin were added to the tube. The tube was incubated  
at 37°C for 1 week. After the incubation, the beads were removed by  
using a centrifugal filter tube (Millipore Co.; UFC30GV00) having  
membrane pores of 0.22 μm. Next, 50 μl of 10% trichloroacetic acid  
was added to the solution (50 μl) from which the beads had been removed.  
20 The mixture was centrifuged to precipitate protein. The protein pellet  
was washed with 300 μl of 5% trichloroacetic acid, and then dried.  
Then, 100 μl of 6 N HCl was added to the protein pellet, and the dissolved  
protein was heated at 110°C for 16 hours. The resulting sample was  
assayed for the quantification of pentosidine by HPLC (T. Miyata et  
25 al., 1996, J. Am. Soc. Nephrol., 7: 1198-1206; T. Miyata et al., 1996,  
Proc. Natl. Acad. Sci. USA, 93: 2353-2358).

The amounts of pentosidine generated by the incubation at 37°C  
are summarized in Fig. 14. It was revealed that the addition of carbonyl  
compound-trapping beads had a strong effect in inhibiting the  
30 generation of pentosidine.

Example 7. Effect of the elimination of carbonyl compounds by carbonyl compound-trapping beads added to the peritoneal dialysate

35 Carbonyl compound-trapping beads were tested for the effect of  
eliminating carbonyl compounds from the peritoneal dialysate.  
Dimethylsulfoxide (100 μl) was added to a tube containing carbonyl

compound-trapping beads (PS-TsNHNH<sub>2</sub>; ARGONAUT TECHNOLOGIES CO.), to swell the beads. Then, 900 µl of a peritoneal dialysate (Baxter Ltd.; Dianeal PD-4,1.5) was added. The mixture was stirred at room temperature for 16 hours by using a rotator. Subsequently, the suspension containing the carbonyl compound-trapping beads was filtered with a centrifugal filter tube (Millipore Co.; UFC30GV00), having membrane pores of 0.22 µm, and the amount of carbonyl compounds in the filtrate was assayed by following method.

<Quantification of carbonyl compounds>

(1) Assay of sample solutions

A sample solution (200 µl) was mixed with 200 µl of a 0.5 N hydrochloric acid solution containing 2,4-DNPH (0.025%), and the mixture was incubated at 30°C for 30 minutes. Subsequently, an aqueous solution of 1 M acetone (20 µl) was added to the mixture. The resulting mixture was incubated at 30°C for 10 minutes. The aqueous mixture was washed 3 times with 200 µl of n-hexane, and 200 µl of octanol was added to the aqueous layer to extract the hydrazone. The octanol layer was recovered and the absorbance thereof was measured at 360 nm in a spectrophotometric microplate reader (Nippon Molecular Devices Co.; SPECTRAMax250).

(2) Preparation of calibration curve

Aqueous solutions of various glucose concentrations were prepared and the amounts of carbonyl compounds derived from glucose were assayed by the same method as described in (i); a calibration curve of glucose concentration vs. concentration of carbonyl compounds was prepared based on this experiment.

(3) Quantification of carbonyl compounds

Glucose concentrations in sample solutions were measured using a glucose assay kit (Wako Pure Chemical Industries, Ltd.; Glucose CII-Test Wako). The amount of carbonyl compounds derived from glucose was estimated by using the calibration curve. The amount of carbonyl compounds in the sample solution was determined by subtracting the amount of glucose-derived carbonyl compounds from the total amount of carbonyl compounds in the sample solution.

The results are shown in Fig. 15. The carbonyl compound-trapping beads (2 mg) were added to the peritoneal dialysate, and the suspension

was stirred at room temperature for 16 hours. This treatment reduced the amount of carbonyl compounds by 55%. When 10 mg of carbonyl compound-trapping beads were added to the dialysate, the amount of carbonyl compounds was further reduced.

5 These findings clarified that the carrier with immobilized carbonyl compound-trapping agent can be used for inhibiting the generation and/or accumulation of carbonyl compounds in the peritoneal dialysate.

10 Example 8. The activity of activated charcoal in trapping carbonyl compounds in a dicarbonyl compound solution

Activated charcoal was used as a carbonyl compound-trapping agent to evaluate its effect of eliminating carbonyl compounds from a dicarbonyl compound solution. A dicarbonyl compound solution was prepared by dissolving a dicarbonyl compound (100  $\mu$ M) in a phosphate buffer (abbreviated hereafter "PBS"). The solution (900  $\mu$ l) was added to a tube containing 25  $\mu$ g or 50  $\mu$ g of activated charcoal (Wako Pure Chemical Industries, Ltd.). Dicarbonyl compounds used were glyoxal, methylglyoxal, and 3-deoxyglucosone. The tube was placed on a rotator and stirred at room temperature for 19 hours. After stirring, the solution in the tube was filtered by a centrifugal filter tube (Millipore Co.; UFC30GV00) having membrane pores of 0.22  $\mu$ m. The concentration of each dicarbonyl compound was determined by high performance liquid chromatography according to a commonly used method.

25 The results are shown in Fig. 16. When 25  $\mu$ g of activated charcoal was added to 900  $\mu$ l of the dicarbonyl compound solution, the activated charcoal trapped glyoxal (GO) by 71%, methylglyoxal (MGO) by 94%, and 3-deoxyglucosone (3DG) by 93%. When 50  $\mu$ g of activated charcoal was used, the charcoal trapped glyoxal by 85%, and both methylglyoxal and 3-deoxyglucosone by 98%. Thus, it was confirmed that most of each dicarbonyl compound tested was trapped by activated charcoal.

30 Example 9. The activity of activated charcoal in trapping carbonyl compound in the peritoneal dialysate

35 Activated charcoal was used as a carbonyl compound-trapping agent to evaluate its effect of eliminating carbonyl compounds from the

peritoneal dialysate. An aliquot (900  $\mu$ l) of peritoneal dialysate (Baxter Ltd.; Dianeal PD-4, 1.5) was added to a tube containing 25  $\mu$ g or 50  $\mu$ g of activated charcoal (Wako Pure Chemical Industries, Ltd.). The tube was placed on a rotator and stirred at room temperature for 19 hours. After stirring, the solution in the tube was filtered by a centrifugal filter tube (Millipore Co.; UFC30GV00) having membrane pores of 0.22  $\mu$ m. The concentrations of glyoxal, methylglyoxal, and 3-deoxyglucosone were determined by high performance liquid chromatography according to a commonly used method.

The results are shown in Fig. 17. When 25  $\mu$ g of activated charcoal was added to 900  $\mu$ l of the peritoneal dialysate, the activated charcoal trapped glyoxal (GO) by 56%, methylglyoxal (MGO) by 71%, and 3-deoxyglucosone (3DG) by 62%, as compared with the dialysates without activated charcoal. When 50  $\mu$ g of activated charcoal was used, the charcoal trapped glyoxal by 64%, methylglyoxal by 78%, and 3-deoxyglucosone by 77%. Thus, it was confirmed that each dicarbonyl compound tested was trapped by activated charcoal.

Example 10. The activities of guanidine, aminoguanidine, and biguanide agents in trapping glyoxal, methylglyoxal, and 3-deoxyglucosone

A mixture (50  $\mu$ l) of glyoxal, methylglyoxal, and 3-deoxyglucosone (1 mM each) was further mixed with 400  $\mu$ l of 0.1 M phosphate buffer (pH 7.4), and each (50  $\mu$ l; 30 mM) of guanidine, aminoguanidine, or a biguanide agent. The resulting mixture was incubated at 37°C. Biguanide agents used were metformin, buformin, and phenformin. After the incubation, glyoxal, methylglyoxal, and 3-deoxyglucosone were converted to quinoxaline derivatives by using o-phenylenediamine, and then the respective concentrations were determined by high performance liquid chromatography.

The results are shown in Fig. 18 (guanidine), Fig. 19 (metformin), Fig. 20 (buformin), Fig. 21 (phenformin), and Fig. 22 (aminoguanidine). Guanidine, aminoguanidine, and all the biguanide agents were shown to have the effect of markedly reducing the concentration of methylglyoxal, in particular. Further, aminoguanidine drastically reduced the concentration of methylglyoxal, and in addition, it has

the effect of markedly reducing the concentration of 3-deoxyglucosone, which was not significantly reduced by other biguanides.

Example 11. The activity of SH agent in trapping glyoxal, methylglyoxal, and 3-deoxyglucosone

A solution (50  $\mu$ l) of glyoxal, methylglyoxal, and 3-deoxyglucosone (1 mM each) was further mixed with 400  $\mu$ l of 0.1 M phosphate buffer (pH 7.4) and an SH compound solution (50  $\mu$ l; 30 mM). The resulting mixture was incubated at 37°C. SH compounds used were cysteine, N-acetylcysteine, and GSH. After the incubation, glyoxal, methylglyoxal, and 3-deoxyglucosone were converted to quinoxaline derivatives by using o-phenylenediamine, and then the respective concentrations were determined by high performance liquid chromatography.

The results are shown in Fig. 23 (cysteine), Fig. 24 (N-acetylcysteine), and Fig. 25 (GSH). All the SH compounds were found to have the effect of markedly reducing the concentrations of both glyoxal and methylglyoxal.

Example 12. The activity of albumin in trapping glyoxal, methylglyoxal, and 3-deoxyglucosone

A solution (50  $\mu$ l) consisting of glyoxal, methylglyoxal, and 3-deoxyglucosone (1 mM each) was further mixed with 400  $\mu$ l of 0.1 M phosphate buffer (pH 7.4), 50  $\mu$ l of 100 mg/ml bovine serum albumin solution, and the resulting mixture was incubated at 37°C. After the incubation, glyoxal, methylglyoxal, and 3-deoxyglucosone were converted to quinoxaline derivatives by using o-phenylenediamine, and then the respective concentrations were determined by high performance liquid chromatography.

The results are shown in Fig. 26. Bovine serum albumin was found to have the effect of markedly reducing the concentrations of glyoxal and methylglyoxal.

Example 13. Effect of SH compounds added to the peritoneal dialysate and incubated at 37°C on inhibition of the generation of pentosidine

An aliquot (490  $\mu$ l) of a peritoneal dialysate (Baxter Ltd.; PD-4,



1.5) was mixed with 70  $\mu$ l of 0.1 M phosphate buffer (pH 7.4) containing an SH compound and 140  $\mu$ l of the peritoneal dialysate (Baxter Ltd.; PD-4, 1.5) containing 30 mg/ml bovine serum albumin. The resulting mixture was incubated at 37°C for 1 week. SH compounds used were cysteine, N-acetylcysteine, and GSH. Aminoguanidine was also used. After the incubation, 50  $\mu$ l of 10% trichloroacetic acid was added to the solution (50  $\mu$ l). The mixture was centrifuged to precipitate the protein. The protein pellet was washed with 300  $\mu$ l of 5% trichloroacetic acid, and then dried. Then, 100  $\mu$ l of 6 N HCl was added to the protein pellet, and the dissolved protein was heated at 110°C for 16 hours. The resulting sample was assayed for the quantification of pentosidine by high performance liquid chromatography (T. Miyata et al., 1996, J. Am. Soc. Nephrol., 7:1198-1206; T. Miyata et al., 1996, Proc. Natl. Acad. Sci. USA., 93:2353-2358).

The results are shown in Fig. 27. It was revealed that the addition of SH compounds had a marked effect of inhibiting the generation of pentosidine.

### Industrial Applicability

The present invention can readily eliminate toxic damages caused by carbonyl compounds during peritoneal dialysis, which had been tormenting patients for a long time. When conventional peritoneal dialysates were used, the peritoneum of a patient being dialyzed was constantly under the carbonyl-stress state caused by carbonyl compounds transferred to the peritoneal cavity from other parts of the body during dialysis, as well as by the carbonyl compounds generated in the process of manufacturing the dialysates. In contrast, the present invention can effectively eliminate carbonyl compounds generated in peritoneal dialysates, and therefore, can sufficiently contribute to the improvement of carbonyl stress of dialysis patients. Furthermore, carbonyl compounds transferred to the peritoneal cavity can also be inactivated or eliminated effectively by infusing a carbonyl compound-trapping agent or by circulating a dialysate through a carbonyl compound-trapping cartridge. Thus, the present invention provides a quite effective approach to prevent damages caused by

peritoneal dialysis including peritoneal damages caused by carbonyl compounds associated with peritoneal dialysis.

In addition, the present invention enables removing carbonyl compounds generated by the degradation of glucose during heat sterilization and long-term storage, thereby successfully providing  
5 a peritoneal dialysate of neutral pH, which has previously been pharmaceutically difficult to prepare because of the degradation of glucose. Thus, the present invention enables a more physiological peritoneal dialysis treatment.

10 The inventive peritoneal dialysate can be practically used with only a simple procedure such as contacting with the carbonyl compound-trapping agent or administering directly. Furthermore, the production of inventive peritoneal dialysate does not require any specific facilities. Thus, the carbonyl compound-trapping agent  
15 based on the present invention, the peritoneal dialysate using the same, and the method of production thereof, creates a new therapeutic concept for peritoneal dialysis treatment.

09762286-082301


## CLAIMS

1. An agent for improving intraperitoneal carbonyl-stress state during peritoneal dialysis, comprising a carbonyl compound-trapping agent as an active ingredient.
2. The agent of claim 1, wherein the carbonyl compound-trapping agent is immobilized on an insoluble carrier.
3. The agent of claim 1, wherein the carbonyl compound-trapping agent is to be mixed with a peritoneal dialysate.
4. The agent of any one of claims 1 to 3, wherein the carbonyl compound-trapping agent is selected from the group consisting of aminoguanidine, pyridoxamine, hydrazine, biguanide compound, SH group containing compound, and derivatives of these.
5. The agent of any one of claims 1 to 3, wherein the carbonyl compound-trapping agent is an agent inhibiting Maillard reaction.
6. The agent of claim 1, wherein the carbonyl compound-trapping agent is a compound insoluble in peritoneal dialysates and capable of adsorbing carbonyl compounds.
7. A cartridge used for trapping carbonyl compounds within peritoneal dialysates, wherein the cartridge is filled with the carbonyl compound-trapping agent(s) of claim 2 and/or claim 6.
8. A method for preparing a peritoneal dialysate having a reduced carbonyl compound content, the method comprising passing the peritoneal dialysate through the cartridge of claim 7.
9. A method for preparing a peritoneal dialysate having a reduced carbonyl compound content, the method comprising:
- (a) contacting the peritoneal dialysate with the carbonyl compound-trapping agent(s) of claim 2 and/or claim 6, and
  - (b) separating the peritoneal dialysate from the carbonyl compound-trapping agent.
10. A peritoneal dialysate comprising a carbonyl compound-trapping agent.
11. The peritoneal dialysate of claim 10, wherein the peritoneal dialysate further comprises a reducing sugar, and is placed in a container comprising a first compartment and a second compartment so that the first compartment contains the reducing sugar and the

second compartment contains the carbonyl compound-trapping agent.  
12. The peritoneal dialysate of claim 10, wherein the carbonyl compound-trapping agent is to be administered into the intraperitoneal cavity.



1/27

Figure 1



2/27


Figure 2



09/763286-082801


3/27

Figure 3



4/27


Figure 4





5/27

Figure 5



6/27

Figure 6






Figure 7




8/27

Figure 8




9/27

Figure 9




10/27

Figure 10




11/27

Figure 11



12/27


Figure 12






13/27

Figure 13



14/27


Figure 14



09/763286-082801


15/27

Figure 15




16/27

Figure 16




17/27

Figure 17




18/27

Figure 18




19/27

Figure 19



20/27


Figure 20






21/27

Figure 21






22/27

Figure 22




23/27

Figure 23




24/27

Figure 24




25/27

Figure 25







26/27

Figure 26



27/27

Figure 27



As a below-named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name,

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

CARBONYL-STRESS IMPROVING AGENT AND PERITONEAL DIALYSATE  
the specification of which (check only one item below):

☐ is attached hereto.

☐ was filed as United States application

Serial No. \_\_\_\_\_

on \_\_\_\_\_

and was amended

on \_\_\_\_\_ (if applicable).

☒ was filed as PCT international application

Number PCT/JP99/04521

on 23 August 1999,

and was amended under PCT Article 19

on \_\_\_\_\_ (if applicable).

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to the examination of this application in accordance with Title 37, Code of Federal Regulations, ' 1.56(a).

I hereby claim foreign priority benefits under Title 35, United States Code, ' 119 of any foreign application(s) for patent or inventor's certificate or of any PCT international application(s) designating at least one country other than the United States of America listed below and have also identified below any foreign application(s) for patent or inventor's certificate or any PCT international application(s) designating at least one country other than the United States of America filed by me on the same subject matter having a filing date before that of the application(s) of which priority is claimed:

## PRIOR FOREIGN/PCT APPLICATION(S) AND ANY PRIORITY CLAIMS UNDER 35 U.S.C. 119:

COUNTRY	APPLICATION	DATE OF FILING	PRIORITY CLAIMED
PCT	PCT/JP99/04521	23 August 1999	<input checked="" type="checkbox"/> YES <input type="checkbox"/> NO
Japan	10/237108	24 August 1998	<input checked="" type="checkbox"/> YES <input type="checkbox"/> NO
Japan	11/155393	02 June 1999	<input checked="" type="checkbox"/> YES <input type="checkbox"/> NO



**Combined Declaration For Patent Application and Power of Attorney (Continued)**  
(Includes Reference to PCT International Applications)

ATTORNEY'S DOCKET  
NUMBER  
SHIM-008

I hereby claim the benefit under Title 35, United States Code, ' 120 of any United States application(s) or PCT international application(s) designating the United States of America that is/are listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in that/those prior application(s) in the manner provided by the first paragraph of Title 35, United States Code, ' 112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, ' 1.56(a) which occurred between the filing date of the prior application(s) and the national or PCT international filing date of this application:

**PRIOR U.S. APPLICATIONS OR PCT INTERNATIONAL APPLICATIONS DESIGNATING THE U.S. FOR BENEFIT UNDER 35 U.S.C. 120:**

U.S. APPLICATIONS

STATUS (Check one)

U.S. APPLICATION NUMBER

U.S. FILING DATE

PATENTED

PENDING

ABANDONED

POWER OF ATTORNEY: As a named inventor, I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application and transact all business in the Patent and Trademark Office connected therewith. (List name and registration number)

Karl Bozicevic, Reg. No. ~~28,807~~  
Bret E. Field, Reg. No. ~~37,620~~  
Dianna L. DeVore, Reg. No. ~~42,484~~  
Alan W. Cannon, Reg. No. ~~34,977~~  
Robert Hall, Reg. No. ~~39,209~~  
Nicole Verona, Reg. No. ~~47,153~~

Carol L. Francis, Reg. No. ~~36,513~~  
Pamela J. Sherwood, Reg. No. ~~36,677~~  
Paula A. Borden, Reg. No. ~~42,344~~  
Carol M. LaSalle, Reg. No. ~~39,740~~  
Kathleen Hall, Reg. No. ~~44,143~~

Send Correspondence to:

**BOZICEVIC, FIELD & FRANCIS LLP**  
200 Middlefield Road, Suite 200  
Menlo Park, CA 94025  
Telephone: (650) 327-3400  
Facsimile: (650) 327-3231

Direct Telephone Calls to:  
(name and telephone number)

Name: Karl Bozicevic  
Registration No. 28,807  
Telephone: (650) 327-3400

FULL NAME OF  
INVENTOR

FAMILY NAME

FIRST GIVEN NAME

SECOND GIVEN NAME

MIYATA

Toshio

RESIDENCE &  
CITIZENSHIP

CITY

STATE OR FOREIGN COUNTRY

COUNTRY OF CITIZENSHIP

Isehara-shi

Kanagawa, Japan

Japan

POST OFFICE  
ADDRESS

POST OFFICE ADDRESS

CITY

STATE & ZIP CODE/COUNTRY

102 Ekuseru Isehara, 16-25, Sakurada  
2-chome,

Isehara-shi

Kanagawa 259-1132 Japan


I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

SIGNATURE OF INVENTOR 201

DATE

F:\DOCUMENTS\SHIM (Shimizu, formerly SHI)\008\SHIM-008 Declaration.doc

United States Patent & Trademark Office  
Office of Initial Patent Examination -- Scanning Division



Application deficiencies found during scanning:

☐ Page(s) \_\_\_\_\_ of \_\_\_\_\_ were not present  
for scanning. (Document title)

☐ Page(s) \_\_\_\_\_ of \_\_\_\_\_ were not  
present  
for scanning. (Document title)

\* Scanned copy is best available. Drawings are done